

a 4-year-old with autism  
was once a 3-year-old  
with autism was once a  
2-year-old with autism  
was once a 1-year-old  
with autism was once a



## Autism can often be recognized at 18 months or younger.

From birth to 5 years, there are milestones children should reach in terms of how they play, learn, speak, and act. A delay in any of these areas could be a sign of a developmental problem, even autism. Fortunately, the earlier a delay is recognized, the more you can do to help them reach their full potential.

**FREE health care professional resource kits and  
additional information are available at  
[www.cdc.gov/actearly](http://www.cdc.gov/actearly) and 1-800-CDC-INFO.**



Learn the Signs. Act Early.

a 4-year-old with autism  
was once a 3-year-old  
with autism was once a  
2-year-old with autism  
was once a 1-year-old  
with autism was once a



## Autism can often be recognized at 18 months or younger.

From birth to 5 years, there are milestones children should reach in terms of how they play, learn, speak, and act. A delay in any of these areas could be a sign of a developmental problem, even autism. Fortunately, the earlier a delay is recognized, the more you can do to help them reach their full potential.

**FREE health care professional resource kits and  
additional information are available at  
[www.cdc.gov/actearly](http://www.cdc.gov/actearly) and 1-800-CDC-INFO.**



Learn the Signs. Act Early.